



PLASTIC SURGERY ASSOCIATES

Account Number: \_\_\_\_\_ Patient Name: \_\_\_\_\_

Address: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Birthdate: \_\_\_\_\_

*Indications – Prescription treatment for hypotrichosis used to grow eyelashes.*

**LATISSE SCREENING INTERVIEW**

- 1. Do you use Lumigan or any other medication to treat intraocular pressure? .....Y .....N
- 2. Are you pregnant or breastfeeding? .....Y .....N
- 3. Do you experience any of the following symptoms / conditions:
  - a. Eye pruritus (itchy eyes) .....Y .....N
  - b. Conjunctival hyperemia (red eyes) .....Y .....N
  - c. Irritated eyes .....Y .....N
  - d. Dry eyes .....Y .....N
  - e. Eyelid redness .....Y .....N
  - f. Twitching eyelids .....Y .....N
  - g. Macular edema .....Y .....N
- 4. Do you have brown spots or discoloration of the eyelids? .....Y .....N

***If you answered Yes to #1 or #2, you may not use Latisse.***

***If you answered Yes to any part of #3, we recommend you see your Ophthalmologist to correct and control your symptoms prior to starting Latisse.***

***If you answered Yes to #4, we recommend you see your Dermatologist before starting Latisse.***

Latisse warnings, precautions and adverse reactions:

- 1. Do not use Latisse with other medications that lower Intraocular Pressure (IOP).
- 2. May cause permanent iris pigmentation if Latisse comes in continual contact with your eye.
- 3. Latisse may cause hyper pigmentation of the eyelid. This is reversible after stopping Latisse.
- 4. It is very important to apply Latisse to skin of upper eyelid margin at the base of the eyelashes to reduce the potential of unwanted hair growth outside treatment area.
- 5. Do not use Latisse if you have Macular edema.
- 6. Avoid contaminating Latisse applicators.
- 7. Do Not use if pregnant or breastfeeding.
- 8. Remove contact lenses prior to applying Latisse. Resume use of contact lenses 15 minutes after Latisse dries.
- 9. Red eye or conjunctal hyperemia will usually resolve in a few days to two (2) weeks. If it persists or suddenly worsens, discontinue Latisse.
- 10. Less than 4% of patients experience eye pruritus (itch), skin hyperpigmentation (darkening), ocular (eye) irritation, dry eye, Erythema (redness) of eyelids.

***If these symptoms do not resolve in a few days, then discontinue Latisse and / or see your Ophthalmologist.***

**\*\*\*Non-refundable – Latisse is a prescription drug - cannot be returned.\*\*\***

Patient Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Witness: \_\_\_\_\_

**Use as directed - see package insert.**